

UNCLASSIFIED

**Defense Technical Information Center  
Compilation Part Notice**

**ADP013402**

**TITLE:** High Dose Pralidoxime [PRX] Treatment Prolongs Time to Extubation [TTE] and Increases Mortality in Paraoxon [POX] Exposed Minipigs

**DISTRIBUTION:** Approved for public release, distribution unlimited

**This paper is part of the following report:**

**TITLE:** Chemical and Biological Medical Treatment Symposium - Industry II World Congress on Chemical and Biological Terrorism

**To order the complete compilation report, use: ADA411272**

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

**The following component part numbers comprise the compilation report:**

**ADP013371 thru ADP013468**

UNCLASSIFIED

### **32. HIGH DOSE PRALIDOXIME (PRX) TREATMENT PROLONGS TIME TO EXTUBATION (TTE) AND INCREASES MORTALITY IN PARAOXON (POX) EXPOSED MINIPIGS**

G. Petroianu, A Missler, K Zuleger, V Ewald, C Thyges, W Bergler, R Ruefer  
Dept. of Pharmacology & Toxicology  
University of Heidelberg at Mannheim  
Maybach St. 14-16;  
68169-Mannheim, Germany

#### **ABSTRACT**

Organophosphates are inhibitors of serine hydrolases. Oximes are clinically available enzyme reactivators.

To determine in vivo the effect of high dose PRX use on "time-to-extubation (TTE)" and mortality as compared to identical therapy without PRX.

12 anaesthetized minipigs were used. All pigs received iv-POX (1mg/kg BW) over 50 minutes. Group one (n = 6) received conventional intensive care therapy as described previously [J Appl Toxicol 18: 293 - 298]. Group two (n = 6) received in addition iv-PRX 10 g ( $\approx$  300 mg/kg BW). Before [base-line;BL], after POX application (50min) and then at 1,2,3,4,8 and 16 hours after POX AChE and BChE activities were measured. Statistics: rank order test; significance for  $p \leq 0.05$ .

In group one TTE was  $7.2 \pm 4.4$  h after last measurement. Mortality was 0. In group two TTE was  $14.7 \pm 6.4$  h after last measurement. Mortality was 4/6.

Pralidoxime therapy has no beneficial effect and its use can not be recommended.

#### **KEYWORDS**

Pralidoxime, reactivators, OP poisoning

(The paper was not presented)